



Northeast Fisheries Science Center Reference Document 15-13

Stock Assessment Update of Summer Flounder for 2015

by Mark Terceiro

Stock Assessment Update of Summer Flounder for 2015

by Mark Terceiro

NOAA Fisheries, Northeast Fisheries Science Center, 166 Water Street, Woods Hole, MA 02543

U.S. DEPARTMENT OF COMMERCE
National Oceanic and Atmospheric Administration
National Marine Fisheries Service
Northeast Fisheries Science Center
Woods Hole, Massachusetts
August 2015

Northeast Fisheries Science Center Reference Documents

This series is a secondary scientific series designed to assure the long-term documentation and to enable the timely transmission of research results by Center and/or non-Center researchers, where such results bear upon the research mission of the Center (see the outside back cover for the mission statement). These documents receive internal scientific review, and most receive copy editing. The National Marine Fisheries Service does not endorse any proprietary material, process, or product mentioned in these documents.

All documents issued in this series since April 2001, and several documents issued prior to that date, have been copublished in both paper and electronic versions. To access the electronic version of a document in this series, go to <http://www.nefsc.noaa.gov/publications/>. The electronic version is available in PDF format to permit printing of a paper copy directly from the Internet. If you do not have Internet access, or if a desired document is one of the pre-April 2001 documents available only in the paper version, you can obtain a paper copy by contacting the senior Center author of the desired document. Refer to the title page of the document for the senior Center author's name and mailing address. If there is no Center author, or if there is corporate (*i.e.*, non-individualized) authorship, then contact the Center's Woods Hole Laboratory Library (166 Water St., Woods Hole, MA 02543-1026).

Information Quality Act Compliance: In accordance with section 515 of Public Law 106-554, the Northeast Fisheries Science Center completed both technical and policy reviews for this report. These predissemination reviews are on file at the NEFSC Editorial Office.

This document may be cited as:

Terceiro M. 2015. Stock assessment update of summer flounder for 2015. US Dept Commer, Northeast Fish Sci Cent Ref Doc. 15-13; 18 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at <http://www.nefsc.noaa.gov/publications/>

TABLE OF CONTENTS

State of Stock	1
Projections.....	2
Catch	3
Stock Distribution and Identification.....	3
Data and Assessment	4
Biological Reference Points (BRPs).....	5
Fishing Mortality	5
Spawning Stock Biomass.....	5
Recruitment.....	5
References Cited.....	6
Figures.....	7

STATE OF STOCK

This assessment of summer flounder (*Paralichthys dentatus*) is an update through 2014 of commercial and recreational fishery catch data, research survey indices of abundance, and the analyses of those data. The summer flounder stock was not overfished but overfishing was occurring in 2014 relative to the biological reference points from the 2013 Stock Assessment Workshop (SAW) 57 benchmark assessment (NEFSC 2013; Figures 1-4). Fishing mortality on the fully selected age 4 fish ranged between 0.793 and 1.776 during 1982-1996 then decreased from 0.867 in 1997 to 0.284 in 2007. Since 2007 the fishing mortality rate has increased to 0.359 in 2014, 16% above the 2013 SAW 57 F_{MSY} proxy = $F_{35\%} = 0.309$ (Figures 1-3). The 90% confidence interval for F in 2014 was 0.274 to 0.435. Spawning stock biomass (SSB) decreased from 24,134 mt in 1982 to 5,394 mt in 1989 and then increased to peaks of 50,357 mt in 2003 and 47,499 mt in 2010. SSB was estimated to be 40,323 mt in 2014, 65% of the 2013 SAW 57 SSB_{MSY} proxy = $SSB_{35\%} = 62,394$ mt, and 29% above the 2013 SAW 57 $\frac{1}{2} SSB_{MSY}$ proxy = $\frac{1}{2} SSB_{35\%} = 31,197$ mt (Figures 1-2, 4). The 90% confidence interval for SSB in 2014 was 35,486 to 49,918 mt. The average recruitment from 1982 to 2014 is 41 million fish at age 0. The 1983 and 1985 year classes are the largest in the assessment time series, at 75 and 62 million fish, while the 1988 year class is the smallest at only 10 million fish. After four below-average year classes in 2010-2013 (36, 20, 23, and 27 million fish), the 2014 year class is currently estimated to be average at 41 million fish (Figures 4-5).

No strong internal model retrospective patterns in F or SSB are evident in the updated assessment model, as the average retrospective errors over the last 7 terminal years are < 15% (Figures 6-7), comparable to the magnitude of the 2013 SAW 57 retrospective errors. The 2014 model estimates of F and SSB adjusted for this minor internal retrospective error are within the model estimate 90% confidence intervals (Figure 2). There is, however, evidence of a recent, consistent, small magnitude pattern in both the underestimation of F and the overestimation of SSB for the last 5 terminal years. There is a stronger and more consistent recent retrospective pattern in recruitment as 5 of the last 7 year classes have been initially over-estimated by a range of 22% to 49% (Figure 8). The historical assessment retrospective likewise indicates the emergence of a gradual upward adjustment of recent F estimates and downward adjustment of recent SSB estimates since the 2011 updated assessment (Figure 9). Tracking of recent assessment estimates and projections of Catch, F , and SSB indicates that while catches have not substantially exceeded the specified Acceptable Biological Catch (ABC; Figure 10), projected F has been exceeded and projected SSB has not been achieved (Figure 11-12). This result is mainly due to the recruitment to the stock of four below-average year classes in 2010-2013.

PROJECTIONS

If the total catch of summer flounder in 2015 equals the specified ABC = 10,329 mt = 22.772 million lbs, the median F in 2015 is projected to be 0.385, above the fishing mortality threshold = F_{MSY} proxy = $F_{35\%}$ = 0.309. The median SSB on November 1, 2015 is projected to be 42,423 mt = 93.527 million lbs, below the biomass target SSB_{MSY} proxy = $SSB_{35\%}$ = 62,394 mt = 137.555 million lbs.

If the stock is fished at the fishing mortality threshold = F_{MSY} proxy = $F_{35\%}$ = 0.309 in 2016-2018, the median annual total catches are the Overfishing Limit (OFL) for 2016-2018. The projected estimates in the following table are medians of the distributions for fixed F.

**OFL Total Catch, Landings, Discards
Fishing Mortality (F) and Spawning Stock Biomass (SSB)
Catches and SSB in metric tons**

Year	Total Catch	Landings	Discards	F	SSB
2015	10,329	8,752	1,577	0.385	42,423
2016	8,194	6,807	1,387	0.309	45,198
2017	8,821	7,304	1,517	0.309	47,900
2018	9,365	8,028	1,337	0.309	50,496

CATCH

Reported 2014 landings in the commercial fishery were 5,134 mt = 11.319 million lbs, about 8% over the commercial quota (4,767 mt = 10.509 million lbs). Estimated 2014 landings in the recreational rod-and-reel fishery were 3,354 mt = 7.394 million lbs, about 6% above the recreational harvest limit (3,179 mt = 7.008 million lbs). Total commercial and recreational landings in 2014 were 8,488 = 18.713 million lbs and total commercial and recreational discard losses were 1,717 mt = 3.785 million lbs, for a total catch in 2014 of 10,205 mt = 22.498 million lbs (Figure 3).

**Catch and Status Table: Summer flounder
(weights in 000s mt, recruitment in millions, arithmetic means)**

Year	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	Max ¹	Min ¹	Mean ¹
Commercial landings	7.7	6.3	4.5	4.1	4.8	6.1	7.5	5.9	5.7	5.1	17.1	4.0	7.5
Commercial discards ³	1.5	1.5	2.1	1.2	1.4	1.5	1.1	0.7	0.7	0.8	2.2	0.2	1.1
Recreational landings	4.9	4.8	4.2	3.7	2.7	2.3	2.6	2.9	3.2	3.4	12.5	1.4	4.7
Recreational discards ³	1.0	0.8	1.0	1.2	1.1	1.1	1.1	0.8	0.7	0.9	1.2	0.1	0.7
Total Catch	14.0	12.1	10.0	9.3	8.8	9.5	11.4	10.3	10.3	10.2	26.3	7.9	13.4
Commercial quota	8.2	6.4	4.7	4.3	5.0	6.0	8.0	5.8	5.4	4.8	8.1	8.2	6.9
Recreational harvest limit	5.5	4.3	3.1	2.9	3.3	4.0	5.3	3.9	3.5	3.2	5.5	2.9	4.6
Spawning Stock Biomass ²	45.4	46.2	44.9	44.8	46.5	47.5	45.4	45.3	41.5	40.3	50.4	5.4	28.8
Recruitment (age 0)	29.5	36.8	38.2	45.2	53.4	36.4	20.3	22.5	27.3	41.4	75.2	9.6	41.8
F (age 4)	0.47	0.35	0.28	0.35	0.35	0.38	0.45	0.36	0.35	0.36	1.78	0.28	0.81

1: Over the period 1982-2014

2: On November 1 annually

3: Dead discards

STOCK DISTRIBUTION AND IDENTIFICATION

The Mid Atlantic Fisheries Management Council (MAFMC) and Atlantic States Marine Fisheries Commission (ASMFC) Fishery Management Plan for summer flounder defines the management unit as all summer flounder from the southern border of North Carolina northeast to the US-Canada border. For assessment purposes, the definition of Wilk et al. (1980) of a unit stock extending from Cape Hatteras north to New England has been accepted in this and previous assessments. The current management unit is consistent with a summer flounder genetics study, which revealed no population subdivision at Cape Hatteras (Jones and Quattro 1999). A consideration of summer flounder stock structure incorporating tagging data supported the existence of stocks north and south of Cape Hatteras, with the stock north of Cape Hatteras

possibly composed of two distinct spawning aggregations, off New Jersey and Virginia-North Carolina (Kraus and Musick 2003). The assessment is consistent with the conclusions of this study.

DATA AND ASSESSMENT

The population model implemented for summer flounder is the forward projecting age-structured model ASAP (Legault and Restrepo 1998; NFT 2013a). The model assumes age-dependent values for instantaneous natural mortality (M) that result in a mean value of $M = 0.25$. The catch in the model includes both commercial and recreational fishery landings and discards at age. The fishery landings and discards are treated as two fleets in the model. Indices of stock abundance including age compositions from the NEFSC winter, spring, and fall; Massachusetts spring and fall; Rhode Island fall and monthly fixed; Connecticut spring and fall; Delaware, New York, New Jersey, VIMS ChesMMAAP, and VIMS NEAMAP spring and fall trawl surveys were used in the ASAP model calibration. Aggregate indices of stock abundance from the URI GSO trawl survey and NEFSC MARMAP and ECOMON larval surveys, and recruitment indices (age 0; Young-Of-the-Year, YOY) from surveys conducted by the states of Massachusetts, Delaware, Maryland, and Virginia were also used in the model calibration.

All of the ongoing research survey indices (expressed as aggregate N) have declined since their most recent peak (generally in 2009-2012) to 2014, in percent as follows:

- NEFSC Spring -49%
- NEFSC Fall -27%
- MADMF Spring -57%
- MADMF Fall -43%
- RIDFW Fall -64%
- RIDFW Monthly -82%
- URIGSO -67%
- CTDEEP Spring -22%
- CTDEEP Fall -54%
- NYDEC -28%
- NJDFW -40%
- DEDFW -17%
- VIMS ChesMMAAP -98%
- NEAMAP Spring -51%
- NEAMAP Fall -60%

Most of the YOY indices suggest good recruitment in 2002, 2004, and 2009, and poorer recruitment since then.

No strong internal model retrospective patterns in F or SSB are evident in the updated assessment model, as the average retrospective errors over the last 7 terminal years are $< 15\%$ (Figures 6-7), comparable to the magnitude of the 2013 SAW 57 retrospective errors. The 2014 model estimates of F and SSB adjusted for this minor internal retrospective error are within the model estimate 90% confidence intervals (Figure 2). There is, however, evidence of a recent, consistent, small-magnitude pattern in both the underestimation of F and the overestimation of SSB for the last 5 terminal years. There is a stronger and more consistent recent retrospective pattern in recruitment as 5 of the last 7 year classes have been initially over-estimated by a range of 22% to 49% (Figure 8). The historical assessment retrospective likewise indicates the

emergence of a gradual upward adjustment of recent F estimates and downward adjustment of recent SSB estimates since the 2011 updated assessment (Figure 9). Tracking of recent assessment estimates and projections of Catch, F, and SSB indicates that while catches have not substantially exceeded the specified ABCs (Figure 10), projected F has been exceeded and projected SSB has not been achieved (Figure 11-12). This result is mainly due to the recruitment to the stock of 4 below average year classes in 2010-2013.

BIOLOGICAL REFERENCE POINTS (BRPS)

The existing 2013 SAW 57 biological reference points for summer flounder are based on stochastic yield and SSB per recruit and stochastic projection models in the NOAA NFT framework (NFT 2013b, c; Thompson and Bell 1934) using values from the 2013 assessment. The fishing mortality reference point is $F_{35\%} = 0.309$ (CV = 15%) as a proxy for F_{MSY} . The biomass reference point proxy is estimated as the projection of January 1, 2013 stock sizes at $F_{35\%} = 0.309$ and mean recruitment of 43 million fish per year (1982-2012). The SSB_{MSY} proxy is estimated to be 62,394 mt (137.6 million lbs; CV = 13%), and the biomass threshold of one-half SSB_{MSY} is estimated to be 31,197 mt (68.8 million lbs; CV = 13%). The MSY proxy is estimated to be 12,945 mt (28.539 million lbs; CV = 13%); 10,455 mt = 23.049 million lbs of landings plus 2,490 mt = 5.490 million lbs of discards).

FISHING MORTALITY

Fishing mortality on the fully selected age 4 fish ranged between 0.793 and 1.776 during 1982-1996 then decreased from 0.867 in 1997 to 0.284 in 2007. Since 2007 the fishing mortality rate has increased to 0.359 in 2014, 16% above the 2013 SAW 57 F_{MSY} proxy = $F_{35\%} = 0.309$ (Figures 1-3). The 90% confidence interval for F in 2014 was 0.274 to 0.435.

SPAWNING STOCK BIOMASS

Spawning stock biomass (SSB) decreased from 24,134 mt in 1982 to 5,394 mt in 1989 and then increased to peaks of 50,357 mt in 2003 and 47,499 mt in 2010. SSB was estimated to be 40,323 mt in 2014, 65% of the 2013 SAW 57 SSB_{MSY} proxy = $SSB_{35\%} = 62,394$ mt, and 29% above the 2013 SAW 57 $\frac{1}{2} SSB_{MSY}$ proxy = $\frac{1}{2} SSB_{35\%} = 31,197$ mt (Figures 1-2 & 4). The 90% confidence interval for SSB in 2014 was 35,486 to 49,918 mt.

RECRUITMENT

The average recruitment from 1982 to 2014 is 41 million fish at age 0. The 1983 and 1985 year classes are the largest in the assessment time series, at 75 and 62 million fish, while the 1988 year class is the smallest at only 10 million fish. After 4 below average year classes in 2010-2013 (36, 20, 23, and 27 million fish), the 2014 year class is currently estimated to be average at 41 million fish (Figures 4-5).

REFERENCES CITED

- Jones WJ, Quattro JM. 1999. Genetic structure of summer flounder (*Paralichthys dentatus*) populations north and south of Cape Hatteras. *Mar Biol.* 133: 129-135.
- Kraus RT, Musick JA. 2003. A brief interpretation of summer flounder, *Paralichthys dentatus*, movements and stock structure with new tagging data on juveniles. *Mar Fish Rev.* 63(3): 1-6.
- Legault CM, Restrepo VR. 1998. A flexible forward age-structured assessment program. *ICCAT Col Vol Sci Pap.* 49:246-253.
- NOAA Fisheries Toolbox (NFT) 2013a. Age Structured Assessment Program (ASAP) version 3.0.11. (Available from: <http://nft.nefsc.noaa.gov>).
- NFT. 2013b. Yield per recruit (YPR) version 3.2.1. (Available from: <http://nft.nefsc.noaa.gov>).
- NFT. 2013c. Age Structured Projection Model (AGEPRO) version 4.2. (Available from: <http://nft.nefsc.noaa.gov>).
- Northeast Fisheries Science Center (NEFSC) 2013. 57th Northeast Regional Stock Assessment Workshop (57th SAW) Assessment Report. US Dept Commerce, Northeast Fish Sci Cent Ref Doc. 13-16, 967 p.
- Thompson WF, Bell FH. 1934. Biological statistics of the Pacific halibut fishery. 2. Effect of changes in intensity upon total yield and yield per unit of gear. *Rep Int Fish (Pacific halibut) Comm.* 8: 49 p.
- Wilk SJ, Smith WG, Ralph DE, Sibunka J. 1980. The population structure of summer flounder between New York and Florida based on linear discriminant analysis. *Trans Am Fish Soc.* 109:265-271.




Figure 1. Stock status time series since 1993 for summer flounder. The horizontal dashed line is the 2013 Stock Assessment Workshop (SAW) 57 fishing mortality threshold reference point proxy; the vertical dashed lines are the 2013 SAW 57 biomass threshold and target reference point proxies.

Summer flounder stock status in 2014




Figure 2. Stock status of summer flounder in 2014 with respect to 2013 Stock Assessment Workshop (SAW) 57 threshold and target reference point proxies. The filled circle is the model estimate and the error bars are 90% confidence intervals; the open circle is the model estimate adjusted for internal model retrospective error.




Figure 3. Total fishery catch and fully-recruited fishing mortality (F, peak at age 4) of summer flounder. The horizontal red line is the 2013 Stock Assessment Workshop (SAW) 57 fishing mortality threshold reference point proxy.




Figure 4. Summer flounder spawning stock biomass (SSB; solid line) and recruitment at age 0 (R; vertical bars) by calendar year. The horizontal dashed line is the 2013 Stock Assessment Workshop (SAW) 57 biomass target reference point proxy; the horizontal red line is the biomass threshold reference point proxy.

Summer flounder S- R Data for 1983-2014 Year Classes




Figure 5. Stock-recruitment scatter plot for summer flounder 1983-2014 year classes.





Figure 6. Results of internal model retrospective analysis for summer flounder: fully recruited F (true age 4, model age 5); average retrospective error = -14%.





Figure 7. Results of internal model retrospective analysis for summer flounder: Spawning Stock Biomass; average retrospective error = +6%.





Figure 8. Results of internal model retrospective analysis for summer flounder: R (recruitment at true age 0, model age 1); average retrospective error = +22%.

Summer Flounder Historical Retrospective 1990-2015 Stock Assessments




Figure 9. Comparison of estimates from the 1990-2015 summer flounder assessments.

Summer flounder Projection Performance: Total Catch




Figure 10. Comparison of estimated and projected catch (ABC) from the 2008 through 2015 summer flounder assessments. The F2015 assessment catches projected for 2016-2018 are the ABCs.

Summer flounder Projection Performance: F




Figure 11. Comparison of estimated and projected fishing mortality (F, peak age 4) from the 2008 through 2015 summer flounder assessments. The F2015 assessment F projected for 2016-2018 corresponds to the projected ABCs.

Summer Flounder Projection Performance: SSB




Figure 12. Comparison of estimated and projected Spawning Stock Biomass (SSB) from the 2008 through 2015 summer flounder assessments. The F2015 assessment SSB projected for 2016-2018 corresponds to the projected ABCs.

Procedures for Issuing Manuscripts in the *Northeast Fisheries Science Center Reference Document (CRD) Series*

Clearance

All manuscripts submitted for issuance as CRDs must have cleared the NEFSC's manuscript/abstract/webpage review process. If any author is not a federal employee, he/she will be required to sign an "NEFSC Release-of-Copyright Form." If your manuscript includes material from another work which has been copyrighted, then you will need to work with the NEFSC's Editorial Office to arrange for permission to use that material by securing release signatures on the "NEFSC Use-of-Copyrighted-Work Permission Form."

For more information, NEFSC authors should see the NEFSC's online publication policy manual, "Manuscript/abstract/webpage preparation, review, and dissemination: NEFSC author's guide to policy, process, and procedure," located in the Publications/Manuscript Review section of the NEFSC intranet page.

Organization

Manuscripts must have an abstract and table of contents, and (if applicable) lists of figures and tables. As much as possible, use traditional scientific manuscript organization for sections: "Introduction," "Study Area" and/or "Experimental Apparatus," "Methods," "Results," "Discussion," "Conclusions," "Acknowledgments," and "Literature/References Cited."

Style

The CRD series is obligated to conform with the style contained in the current edition of the United States Government Printing Office Style Manual. That style manual is silent on many aspects of scientific manuscripts. The CRD series relies more on the CSE Style Manual. Manuscripts should be prepared to conform with these style manuals.

The CRD series uses the American Fisheries Society's guides to names of fishes, mollusks, and decapod

crustaceans, the Society for Marine Mammalogy's guide to names of marine mammals, the Biosciences Information Service's guide to serial title abbreviations, and the ISO's (International Standardization Organization) guide to statistical terms.

For in-text citation, use the name-date system. A special effort should be made to ensure that all necessary bibliographic information is included in the list of cited works. Personal communications must include date, full name, and full mailing address of the contact.

Preparation

Once your document has cleared the review process, the Editorial Office will contact you with publication needs – for example, revised text (if necessary) and separate digital figures and tables if they are embedded in the document. Materials may be submitted to the Editorial Office as files on zip disks or CDs, email attachments, or intranet downloads. Text files should be in Microsoft Word, tables may be in Word or Excel, and graphics files may be in a variety of formats (JPG, GIF, Excel, PowerPoint, etc.).

Production and Distribution

The Editorial Office will perform a copy-edit of the document and may request further revisions. The Editorial Office will develop the inside and outside front covers, the inside and outside back covers, and the title and bibliographic control pages of the document.

Once both the PDF (print) and Web versions of the CRD are ready, the Editorial Office will contact you to review both versions and submit corrections or changes before the document is posted online.

A number of organizations and individuals in the Northeast Region will be notified by e-mail of the availability of the document online.

Research Communications Branch
Northeast Fisheries Science Center
National Marine Fisheries Service, NOAA
166 Water St.
Woods Hole, MA 02543-1026

**MEDIA
MAIL**

Publications and Reports of the Northeast Fisheries Science Center

The mission of NOAA's National Marine Fisheries Service (NMFS) is "stewardship of living marine resources for the benefit of the nation through their science-based conservation and management and promotion of the health of their environment." As the research arm of the NMFS's Northeast Region, the Northeast Fisheries Science Center (NEFSC) supports the NMFS mission by "conducting ecosystem-based research and assessments of living marine resources, with a focus on the Northeast Shelf, to promote the recovery and long-term sustainability of these resources and to generate social and economic opportunities and benefits from their use." Results of NEFSC research are largely reported in primary scientific media (*e.g.*, anonymously-peer-reviewed scientific journals). However, to assist itself in providing data, information, and advice to its constituents, the NEFSC occasionally releases its results in its own media. Currently, there are three such media:

NOAA Technical Memorandum NMFS-NE -- This series is issued irregularly. The series typically includes: data reports of long-term field or lab studies of important species or habitats; synthesis reports for important species or habitats; annual reports of overall assessment or monitoring programs; manuals describing program-wide surveying or experimental techniques; literature surveys of important species or habitat topics; proceedings and collected papers of scientific meetings; and indexed and/or annotated bibliographies. All issues receive internal scientific review and most issues receive technical and copy editing.

Northeast Fisheries Science Center Reference Document -- This series is issued irregularly. The series typically includes: data reports on field and lab studies; progress reports on experiments, monitoring, and assessments; background papers for, collected abstracts of, and/or summary reports of scientific meetings; and simple bibliographies. Issues receive internal scientific review and most issues receive copy editing.

Resource Survey Report (formerly *Fishermen's Report*) -- This information report is a regularly-issued, quick-turnaround report on the distribution and relative abundance of selected living marine resources as derived from each of the NEFSC's periodic research vessel surveys of the Northeast's continental shelf. This report undergoes internal review, but receives no technical or copy editing.

TO OBTAIN A COPY of a *NOAA Technical Memorandum NMFS-NE* or a *Northeast Fisheries Science Center Reference Document*, either contact the NEFSC Editorial Office (166 Water St., Woods Hole, MA 02543-1026; 508-495-2350) or consult the NEFSC webpage on "Reports and Publications" (<http://www.nefsc.noaa.gov/nefsc/publications/>). To access *Resource Survey Report*, consult the Ecosystem Surveys Branch webpage (<http://www.nefsc.noaa.gov/femad/ecosurvey/mainpage/>).

ANY USE OF TRADE OR BRAND NAMES IN ANY NEFSC PUBLICATION OR REPORT DOES NOT IMPLY ENDORSEMENT.